



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES
(Untabulated).

ALGERIA—*Oran*.—Month of February, 1912. Population, 123,086. Total number of deaths from all causes 191, including diphtheria 2, measles 7, tuberculosis 29, typhoid fever 3.

BRAZIL—*Pernambuco*.—Two weeks ended January 31, 1912. Population, 225,000. Total number of deaths from all causes 392, including plague 2, smallpox 64, tuberculosis 60, yellow fever 1.

BRITISH EAST AFRICA—*Mombasa*.—Month of January, 1912. Population, 26,000. Total number of deaths from all causes 55, including tuberculosis 6.

FRANCE—*Marseille*.—Month of February, 1912. Population, 550,619. Total number of deaths from all causes 806, including diphtheria 6, measles 1, smallpox 1, tuberculosis 126, typhoid fever 8.

Roubaix.—Month of February, 1912. Population, 122,723. Total number of deaths from all causes 168, including scarlet fever 2, tuberculosis 17, typhoid fever 1.

GREAT BRITAIN.—Week ended March 2, 1912:

England.—The deaths registered in 77 great towns correspond to an annual rate of 15.4 per 1,000 of the population, which is estimated at 17,559,219.

Ireland.—The deaths registered in 21 principal town districts correspond to an annual rate of 22.1 per 1,000 of the population, which is estimated at 1,157,014. The lowest rate was recorded at Sligo, viz, 4.4, and the highest at Newtownards, viz, 40.1.

Scotland.—The deaths registered in 18 principal towns correspond to an annual rate of 18.7 per 1,000 of the population, which is estimated at 2,182,400. The lowest rate was recorded at Kirkaldy, viz, 10.3, and the highest in Greenock, viz, 24. The total number of deaths from all causes was 784, including diphtheria 9, measles 56, scarlet fever 6, typhoid fever 1.

JAMAICA—*Kingston*.—Month of February, 1912. Population, 52,000. Total number of deaths from all causes 157, including tuberculosis, pulmonary, 26, typhoid fever 4.

NEW ZEALAND.—Month of November, 1911.

Auckland.—Population, 102,676. Total number of deaths 73, including tuberculosis 3.

Christchurch.—Population, 80,193. Total number of deaths 45, including diphtheria 1, tuberculosis 3.

Dunedin.—Population, 64,237. Total number of deaths 54, including tuberculosis 5.

Wellington.—Population, 70,729. Total number of deaths 57, including diphtheria 1, tuberculosis 5.

ST. THOMAS.—Three months ended December 31, 1911. Population, 11,912. Total number of deaths from all causes 202, including diphtheria 2, tuberculosis, pulmonary, 4, typhoid fever 1.

SIAM—*Bangkok*.—Four weeks ended January 27, 1912. Population, 600,000. Total number of deaths from all causes not reported. The deaths include cholera 196, plague 1, smallpox 700.

SIERRA LEONE—*Freetown*.—Month of January, 1912. Population, 40,000. Total number of deaths from all causes 79. No contagious diseases.

SOUTH AFRICA—*Johannesburg*.—Two weeks ended February 10, 1912. Population, 237,220. Total number of deaths from all causes 140, including diphtheria 1, measles 1, scarlet fever 1, tuberculosis 31, typhoid fever 14.

VENEZUELA—*La Guaira*.—Two weeks ended February 29, 1912. Population, 10,000. Total number of deaths from all causes 21 including tuberculosis 1, yellow fever 1.

By authority of the Secretary of the Treasury:

RUPERT BLUE,
Surgeon General,

United States Public Health and Marine-Hospital Service.

ADDITIONAL COPIES of this publication
may be procured from the SUPERINTEND-
ENT OF DOCUMENTS, Government Printing
Office, Washington, D. C., at 5 cents per copy
Subscription price, per volume..... \$2

